

ProCalc Practice Questions: Question Type 8 (dosage/dose or day BSA)

- 1.** Cytosar-U (Cytarabine) has been prescribed for a patient with a current BSA of 0.5. The drug literature recommends 100-200 mg/m²/day. In checking the appropriateness of the drug order, you determine that ____ to ____ mg of Cytosar-U would be an appropriate dose per day.
- 2.** Neomycin has been prescribed for a patient with a current BSA of 0.54. The drug literature recommends 2-7 g/m²/day. In checking the appropriateness of the drug order, you determine that ____ to ____ g of Neomycin would be an appropriate dose per day.
- 3.** Cytosar-U (Cytarabine) has been prescribed for a patient with a current BSA of 0.56. The drug literature recommends 100-200 mg/m²/day. In checking the appropriateness of the drug order, you determine that ____ to ____ mg of Cytosar-U would be an appropriate dose per day.
- 4.** Vepesid (Etoposide) has been prescribed for a patient with a current BSA of 2. The drug literature recommends 35 mg/m²/day. In checking the appropriateness of the drug order, you determine that ____ mg of Vepesid would ____ be an appropriate dose per day.
- 5.** You need to verify that a prescribed dose of Leucovorin for a patient with a current BSA of 1.82 is appropriate. The drug literature recommends 10mg/m²/dose. You determine that an appropriate dose for this patient would be ____ mg per dose.

ProCalc Practice Questions: Question Type 8 ANSWERS

- 1.** 50; 100
- 2.** 1.1;3.81
- 3.** 56; 112
- 4.** 70
- 5.** 18